

Table 6-18
TRVs Selected for Benthic Invertebrate Tissue Assessment

Constituent	Units	Benthic Invertebrates - Tissue TRVs						
		NBSA BERA TRV			TRVs Developed Specifically for NBSA BERA	LPR FFS TRV (USEPA 2014)		
		NOAEL	LOAEL	Source		NOAEL	LOAEL	Source
Dioxins/Furans								
2,3,7,8-TCDD	mg/kg, ww	0.0003 ^a	0.003	Crayfish; survival (Ashley et al. 1996)		0.00000015	0.0000013	Eastern oyster; reproduction (Wintermeyer and Cooper 2003)
Polychlorinated Biphenyls								
Total PCBs	mg/kg, ww	0.052 ^a	0.52	10 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.0064	0.017	reproduction eastern oyster (Chu et al 2000; 2003)
Pesticides								
Total DDx (2,4 & 4,4)	mg/kg, ww	0.011 ^a	0.11	6 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.06	0.13	Pink shrimp; mortality (Nimmo et al. 1970)
Total DDx (2,4 & 4,4) Alternative TRV	mg/kg, ww	0.001 ^a	0.01	An alternative distribution (beta general) was selected to derive an alternative 5th percentile LOAEL TRV (10 µg/kg ww) as a conservative SSD-derived estimate. An alternative NOAEL (1.0 µg/kg ww) was extrapolated from the alternative 5th percentile LOAEL TRV using an uncertainty factor of 10.		--	--	--
Dieldrin	mg/kg, ww	0.008 ^a	0.08	Pink shrimp; survival (Parrish et al. 1973)		0.0016	0.008	Pink shrimp; survival (Parrish et al. 1973)
Total Chlordane	mg/kg, ww	0.71	1.7	Pink shrimp; mortality (Parrish et al. 1976)	X	--	--	--
Hexachlorobenzene	mg/kg, ww	10.6	15.8	Amphipod; mortality (Nebeker et al. 1989)	X	--	--	--
Polycyclic Aromatic Hydrocarbons								
Total LMW PAH	mg/kg, ww	11 ^a	111	Amphipod; mortality (Lee et al. 2002; fluorene study)		0.078	0.78	Polychaete worm; reproduction (Emery and Dillon 1996)
Total HMW PAH	mg/kg, ww	8.1	22.2	Amphipod; growth and reproduction (Schuler et al. 2007; fluoranthene study)		0.066	0.66	Blue mussel; reproduction (Eertman et al. 1995)
Inorganics								
Arsenic	mg/kg, ww	0.064 ^a	0.64	7 species; survival and growth. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		--	--	--
Cadmium	mg/kg, ww	0.024 ^a	0.24	29 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		--	--	--
Chromium	mg/kg, ww	1.5	3.5	Amphipod; survival and growth (Norwood et al. 2007)		--	--	--
Copper	mg/kg, ww	--	--	--		5	12	Baltic clam; survival (Absil et al. 1996)
Lead	mg/kg, ww	4 ^a	40	Amphipod; survival (Spehar et al. 1978)		0.52	2.6	Amphipod (Hyalalela); survival (Borgmann and Norwood 1999)
Mercury/Methylmercury	mg/kg, ww	0.048	0.095	Whole body copepod; reproduction (Hook and Fisher 2002)		0.048	0.095	Whole body copepod; reproduction (Hook and Fisher 2002)
Nickel	mg/kg, ww	0.1 ^a	1.1	Amphipod; survival (Borgmann et al. 2001)				
Selenium	mg/kg, ww	0.05	0.51	Midge; growth (Malchow et al. 1995)		--	--	--
Silver	mg/kg, ww	0.49	0.59	Water flea; growth and reproduction (Naddy et al. 2007)		--	--	--
Zinc	mg/kg, ww	5.1 ^a	51	Crustaceans; survival (Muyssen et al. 2006		--	--	--

Table 6-18
TRVs Selected for Benthic Invertebrate Tissue Assessment

Notes:

^a NOAEL extrapolated from LOAEL using an uncertainty factor of 10.

Acronyms and Abbreviations:

"--" = not available

BERA = baseline ecological risk assessment

FFS = focused feasibility study

HMW = high molecular weight

LMW = low molecular weight

LOAEL = lowest observed adverse effects level

LPR = Lower Passaic River

mg/kg, ww = milligrams per kilogram, wet weight

NBSA = Newark Bay Study Area

NOAEL = no observed adverse effect level

PAH = polycyclic aromatic hydrocarbon

PCB = polychlorinated biphenyl

SSD = species sensitivity distribution

TCDD = tetrachlorodibenzo-p-dioxin

Total DDx = sum of dichlorodiphenyldichloroethylene (DDE), dichlorodiphenyldichloroethane (DDD), and dichlorodiphenyltrichloroethane (DDT)

TRV = toxicity reference value

References:

Absil, M.C.P., M. Berntssen, and L. Gerringa. 1996. The influence of sediment, food and organic ligands on the uptake of copper by sediment-dwelling bivalves. *Aquat Toxicol* 34:13-29.

Ashley, C.M., M.G. Simpson, D.M. Holdich, and D.R. Bell. 1996. 2,3,7,8-tetrachloro-dibenzo-p-dioxin is a potent toxin and induces cytochrome P450 in the crayfish, *Pacifastacus leniusculus*. *Aquat Toxicol* 35:157-69.

Borgmann, U., R. Neron, and W.P. Norwood. 2001. Quantification of bioavailable nickel in sediments and toxic thresholds to *Hyalella azteca*. *Environ Pollut* 111:189-198

Chu, F-LE, P. Soudant, and R.C. Hale. 2003. Relationship between PCB accumulation and reproductive output in conditioned oysters *Crassostrea virginica* fed a contaminated algal diet. *Aqua Tox* 65:293-307.

Chu, F-LE, P. Soudant, L.A. Cruz-Rodriguez, and R.C. Hale. 2000. PCB uptake and accumulation by oysters (*Crassostrea virginica*) exposed via a contaminated algal diet. *Mar Environ Res* 50:217-221.

Eertman, R.H.M., C.L. Groenink, B. Sandee, H. Hummel, and A.C Small. 1995. Response of the blue mussel *Mytilus edulis* L. following exposure to PAHs or contaminated sediment. *Mar Environ Res* 39:169-173.

Emery, V.L. and T.M. Dillon. 1996. Chronic toxicity of phenanthrene to the marine polychaete worm, *Nereis* (Neanthes) arenaceodentata. *Bull Environ Contam Toxicol* 56(2):265-270.

Hansen, D.J., P.R. Parrish, and J. Forester. 1974. Aroclor 1016: Toxicity to and uptake by estuarine animals. *Environ Res* 7:363-373.

Hook, S.E. and N.S. Fisher. 2002. Relating the reproductive toxicity of five ingested metals in calanoid copepods with sulfur affinity. *Mar Environ Res* 53:161-174.

Lee J-H, Landrum P.F., and C-H Koh. 2002. Toxicokinetics and time-dependent PAH toxicity in the amphipod *Hyalella azteca*. *Environ Sci Technol* 36:3124-3130.

Malchow, D.E., A.W. Knight, and K.J. Maier. 1995. Bioaccumulation and toxicity of selenium in *Chironomus decorus* larvae fed a diet of seleniferous *Selenastrum capricornutum*. *Arch Environ Contam Toxicol* 29:104-109

Muyssen BTA, De Schampelaere KAC, Janssen CR. 2006. Mechanisms of chronic waterborne Zn toxicity in *Daphnia magna*. *Aquat Toxicol* 77:393-401.

Naddy, R.B., J.W. Gorsuch, A.B. Rehner, G.R. Mc Nerney, R.A. Bell, and J.R. Kramer. 2007. Chronic toxicity of silver nitrate to *Ceriodaphnia dubia* and *Daphnia magna*, and potential mitigating factors. *Aquatic Toxicology* 84(2):I-X.

Nebeker, A.V., W.L. Griffis, C.M. Wise, E. Hopkins, and J.A. Barbita. 1989. Survival, reproduction and bioconcentration in invertebrates and fish exposed to hexachlorobenzene. *Environ Toxicol Chem* 8:601-11.

Nimmo, D.R., A.J. Wilson Jr., and R.R. Blackman. 1970. Localization of DDT in the body organs of pink and white shrimp. *Bull Environ Contam Toxicol* 5(4):333-341.

Norwood, W.P., U. Borgmann, and D.G. Dixon. 2007. Chronic toxicity of arsenic, cobalt, chromium and manganese to *Hyalella azteca* in relation to exposure and bioaccumulation. *Environ Pollut* 147:262-272.

Parrish, P.R., J.A. Couch, J. Forester, J.M. Patrick, Jr., and G.H. Cook. 1973. Dieldrin: effects on several estuarine organisms. Contrib. No. 178 Gulf Breeze Environ. Res. Lab. In Proceeding of the 27th An. Conf. of the Southeastern Assoc, of Game and Fish. Commissioners.

Parrish, P.R., S.C. Schimmel, D.J. Hansen, J.M. Patrick, Jr., and J. Forester. 1976. Chlordane: effects on several estuarine organisms. *J Toxicol Environ Health* 1:485-494.

Schuler L.J., Landrum P.F., and M.J. Lydy. 2007. Response spectrum of pentachlorobenzene and fluoranthene for *Chironomus tentans* and *Hyalella azteca*. *Environ Toxicol Chem* 26(6):1248-1257.

Sofyan A, Rosita G, Price D, Birge WJ. 2007. Cadmium uptake by *Ceriodaphnia dubia* from different exposures: relevance to body burden and toxicity. *Environ Toxicol Chem* 26(3):470-477.

Spehar RL, Anderson RL, Fiandt JT. 1978. Toxicity and bioaccumulation of cadmium and lead in aquatic invertebrates. *Environ Pollut* 15:195-208.

USEPA. 2014. Focused feasibility study report for the lower eight miles of the Lower Passaic River. The Louis Berger Group, Battelle, and HDR/HydroQual.

Windward Environmental LLC. 2019. Lower Passaic River Study Area Baseline Ecological Risk Assessment. Final. June

Wintermyer ML, Cooper KR. 2003. Dioxin/furan and polychlorinated biphenyl concentrations in eastern oyster (*Crassostrea virginica* Gmelin) tissues and the effects on egg fertilization and development. *J Shellfish Res* 22(3):737-746

Table 6-19a
Calculated Tissue HQs for Benthic Invertebrates with NBSA- and LPR FFS-Selected TRVs (NBSA-wide)

Constituent	Units	Benthic Invertebrate Tissue - HQs (NBSA-wide)											
		Polychaete Worm				Blue Crab				Softshell Clam			
		NBSA TRV-based HQ		LPR FFS TRV-based HQ		NBSA TRV-based HQ		LPR FFS TRV-based HQ		NBSA TRV-based HQ		LPR FFS TRV-based HQ	
		NOAEL	LOAEL	NOAEL	LOAEL	NOAEL	LOAEL	NOAEL	LOAEL	NOAEL	LOAEL	NOAEL	LOAEL
Dioxins/Furans													
2,3,7,8-TCDD	unitless	0.04	0.004	75	9	0.08	0.008	167	19	0.02	0.002	44	5
Polychlorinated Biphenyls													
Total PCBs	unitless	1	0.1	10	4	7	0.7	57	21	3	0.3	22	8
Pesticides													
Total DDX (2,4 & 4,4)	unitless	0.4	0.04	0.07	0.03	11	1	2	0.9	2	0.2	0.4	0.2
Total DDX (2,4 & 4,4) Alternative TRV	unitless	4	0.4	NC	NC	117	12	NC	NC	26	3	NC	NC
Dieldrin	unitless	0.3	0.03	1	0.3	0.7	0.07	4	0.7	0.08	0.008	0.4	0.08
Total Chlordane	unitless	0.003	0.001	NC	NC	0.05	0.02	NC	NC	0.005	0.002	NC	NC
Hexachlorobenzene	unitless	0.00004	0.00002	NC	NC	0.0001	0.00007	NC	NC	0.00003	0.00002	NC	NC
Polycyclic Aromatic Hydrocarbons													
Total HMW PAH	unitless	0.006	0.002	0.8	0.08	0.005	0.002	0.6	0.06	0.05	0.02	6	0.6
Total LMW PAH	unitless	0.001	0.0001	0.2	0.02	0.006	0.0006	0.9	0.09	0.005	0.0005	0.8	0.08
Inorganics													
Arsenic	unitless	43	4	NC	NC	41	4	NC	NC	92	9	NC	NC
Cadmium	unitless	4	0.4	NC	NC	15	1	NC	NC	9	0.9	NC	NC
Chromium	unitless	0.4	0.2	NC	NC	0.2	0.08	NC	NC	3	1	NC	NC
Copper	unitless	NC	NC	0.4	0.2	NC	NC	6	3	NC	NC	3	1
Lead	unitless	0.2	0.02	1	0.3	0.3	0.03	2	0.5	7	0.7	56	11
Mercury	unitless	0.6	0.3	0.6	0.3	3	1	3	1	4	2	4	2
Methylmercury	unitless	0.08	0.04	0.08	0.04	3	1	3	1	0.9	0.4	0.9	0.4
Nickel	unitless	5	0.5	NC	NC	3	0.3	NC	NC	28	3	NC	NC
Selenium	unitless	7	0.7	NC	NC	22	2	NC	NC	19	2	NC	NC
Silver	unitless	0.08	0.06	NC	NC	2	1	NC	NC	0.9	0.7	NC	NC
Zinc	unitless	7	0.7	NC	NC	8	0.8	NC	NC	9	0.9	NC	NC

Table 6-19a**Calculated Tissue HQs for Benthic Invertebrates with NBSA- and LPR FFS-Selected TRVs (NBSA-wide)****Notes:**

Shaded/bolded HQ values indicate an HQ greater than 1.

HQs are calculated using an exposure point concentration based on the 95 percent upper confidence limit, except in the following instances:

- If the number of detected samples or sample size was less than 6, the maximum detected concentration was used.
- If the number of detected samples was 0, the maximum non-detect concentration was used.

Acronyms and Abbreviations:

FFS = focused feasibility study (USEPA 2014)

HMW = high molecular weight

HQ = hazard quotient

LMW = low molecular weight

LOAEL = lowest observed adverse effect level

LPR = Lower Passaic River

NBSA = Newark Bay Study Area

NC = not calculated; TRV not available

NOAEL = no observed adverse effect level

PAH = polycyclic aromatic hydrocarbon

PCB = polychlorinated biphenyl

TCDD = tetrachlorodibenzo-p-dioxin

Total DDx = sum of dichlorodiphenyldichloroethylene (DDE), dichlorodiphenyldichloroethane (DDD), and dichlorodiphenyltrichloroethane (DDT)

TRV = toxicity reference value

References:

USEPA. 2014. Focused feasibility study report for the lower eight miles of the Lower Passaic River. The Louis Berger Group, Battelle, and HDR/HydroQual.

Table 7-5
TRVs for Fish Tissue Assessment

Constituent	Units	Fish - Tissue TRVs						
		NBSA BERA TRV			TRVs Developed Specifically for NBSA BERA	LPR FFS TRV (USEPA 2014)		
		NOAEL	LOAEL	Source		NOAEL	LOAEL	Source
Dioxins/Furans								
2,3,7,8-TCDD	mg/kg, ww	0.000012 ^a	0.00012	7 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.00000089	0.0000018	Mummichog; prey capture behavior (Couillard et al. 2011)
Total Dioxin/Furan TEQ Fish	mg/kg, ww	0.000012 ^a	0.00012	7 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.00000089	0.0000018	Mummichog; prey capture behavior (Couillard et al. 2011)
Total Dioxin/Furan TEQ Fish Alternative TRV	mg/kg, ww	0.0000023	0.000023	As an alternative, a 5th percentile LOAEL based on the beta general distribution (2.3E-05 mg/kg ww)—which most accurately predicts the lowest LOAEL but has a relatively poor visual and statistical fit to the empirical data (when compared with the other distributions noted above)—was selected as a conservative SSD-derived estimate. The alternative NOAEL TRV (2.3E-06 mg/kg ww) was extrapolated from the LOAEL TRV using an uncertainty factor of 10.		--	--	--
Polychlorinated Biphenyls								
Total PCBs	mg/kg, ww	0.38 ^a	3.8	11 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.17	0.53	Atlantic salmon smolt seawater preference; behavior (Lerner et al. 2007)
Total PCB TEQ Fish	mg/kg, ww	0.000012 ^a	0.00012	7 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.00000089	0.0000018	Mummichog; prey capture behavior (Couillard et al. 2011)
Total PCB TEQ Fish Alternative TRV	mg/kg, ww	0.0000023	0.000023	As an alternative, a 5th percentile LOAEL based on the beta general distribution (2.3E-05 mg/kg ww)—which most accurately predicts the lowest LOAEL but has a relatively poor visual and statistical fit to the empirical data (when compared with the other distributions noted above)—was selected as a conservative SSD-derived estimate. The alternative NOAEL TRV (2.3E-06 mg/kg ww) was extrapolated from the LOAEL TRV using an uncertainty factor of 10.		--	--	--
Total Dioxin/Furan/PCB TEQ Fish	mg/kg, ww	0.000012 ^a	0.00012	7 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.00000089	0.0000018	Mummichog; prey capture behavior (Couillard et al. 2011)
Total Dioxin/Furan/PCB TEQ Fish Alternative TRV	mg/kg, ww	0.0000023	0.000023	As an alternative, a 5th percentile LOAEL based on the beta general distribution (2.3E-05 mg/kg ww)—which most accurately predicts the lowest LOAEL but has a relatively poor visual and statistical fit to the empirical data (when compared with the other distributions noted above)—was selected as a conservative SSD-derived estimate. The alternative NOAEL TRV (2.3E-06 mg/kg ww) was extrapolated from the LOAEL TRV using an uncertainty factor of 10.		--	--	--
Pesticides								
Total DDx (2,4 & 4,4)	mg/kg, ww	0.052 ^a	0.52	7 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.078	0.39	9 species; growth, survival, reproduction, and behavior (Beckvar et al. 2005)
Dieldrin	mg/kg, ww	0.12	0.2	Rainbow trout; survival (Shubat and Curtis 1986)		0.008	0.04	Rainbow trout; survival (Shubat and Curtis 1986)
Total Chlordane	mg/kg, ww	1.66 ^a	16.6	Pinfish; mortality (Parrish et al. 1976)	X	--	--	--
Hexachlorobenzene	mg/kg, ww	468	Not derived	Fathead minnow; mortality (Schuytema et al. 1990)	X	--	--	--

Table 7-5
TRVs for Fish Tissue Assessment

Constituent	Units	Fish - Tissue TRVs						
		NBSA BERA TRV			TRVs Developed Specifically for NBSA BERA	LPR FFS TRV (USEPA 2014)		
		NOAEL	LOAEL	Source		NOAEL	LOAEL	Source
Inorganics								
Arsenic	mg/kg, ww	1.3	2.5	Rainbow trout; growth (Erickson et al. 2011)		--	--	--
Chromium	mg/kg, ww	1.28	1.3	Chinook salmon; mortality (Farag et al. 2006)	X	--	--	--
Copper	mg/kg, ww	3.92	4.48	Rainbow trout; mortality (Mount et al. 1994)	X	0.32	1.5	Striped mullet; survival (Zyadah and Abdel-Baky 2000)
Lead	mg/kg, ww	2.5	4.0	Brook trout; growth (Holcombe et al. 1976)		0.4 ^a	4.0	Brook trout; reproduction (Holcombe et al. 1976)
Mercury/Methylmercury	mg/kg, ww	0.035 ^a	0.35	12 species; survival, growth, reproduction, and behavior. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.052	0.26	7 species; growth, survival, reproduction, and behavior (Beckvar et al. 2005)
Selenium	mg/kg, ww	N/A ^b	1.6	Bluegill, sunfish, fathead minnow; reproduction (Coyle et al. 1993; Hermanutz et al. 1992; Ogle and Knight 1989)		--	--	--
Silver	mg/kg, ww	0.11	0.24	Rainbow trout; growth (Guadagnolo et al. 2001)		--	--	--
Zinc	mg/kg, ww	287	403	Guppy; growth (Pearson 1981)		--	--	--

Notes

- ^a NOAEL extrapolated from LOAEL using an uncertainty factor of 10.
- ^b No NOAEL was selected because LOAEL based on ED10 value for the most sensitive species evaluated, below which adverse effects are not expected.

Acronyms and Abbreviations:

"--" = not available	NOAEL = no observed adverse effect level
BERA = baseline ecological risk assessment	PCB = polychlorinated biphenyl
FFS = focused feasibility study	SSD = species sensitivity distribution
LOAEL = lowest observed adverse effects level	TCDD = tetrachlorodibenzo-p-dioxin
LPR = Lower Passaic River	TEQ = toxic equivalent
mg/kg, ww = milligrams per kilogram, wet weight	Total DDx = sum of dichlorodiphenyldichloroethylene (DDE), dichlorodiphenyldichloroethane (DDD), and dichlorodiphenyltrichloroethane (DDT)
N/A = Not applicable; no NOAEL was selected because LOAEL based on ED10 value for the most sensitive species evaluated, below which adverse effects are not expected	TRV = toxicity reference value
NBSA = Newark Bay Study Area	

References:

Beckvar, N., T.M. Dillon, and L.B. Read. 2005. Approaches for linking whole-body fish tissue residues of mercury or DDT to biological effects thresholds. *Environ Toxicol Chem* 24(8):2094-2105.

Couillard, C.M., B. Legare, A. Bernier, and Z. Dionne. 2011. Embryonic exposure to environmentally relevant concentrations of PCB126 affects prey capture ability of *Fundulus heteroclitus* larvae. *Mar Environ Res* 71(4).

Coyle JJ, Buckler DR, Ingersoll CG, Fairchild JF, May TW. 1993. Effect of dietary selenium on the reproductive success of bluegills (*Lepomis macrochirus*). *Environ Toxicol Chem* 12:551-565.

Erickson, R.J., D.R. Mount, T.L. Highland, J.R. Hockett, and C.T. Jenson. 2011. The relative importance of waterborne and dietborne arsenic exposure on survival and growth of juvenile rainbow trout. *Aquatic Toxicology* 104:108-115.

Farag, A.M., T. May, G.D. Marty, M.D.L. Easton, D.D. Harper, E.E. Little, and L. Cleveland. 2006. The effect of chronic chromium exposure on the health of Chinook salmon (*Oncorhynchus tshawytscha*). *Aquat Toxicol* 76:246-257.

Fisk AT, Yarechewski AL, Metner DA, Evans RE, Lockhart WL, Muir DCG. 1997. Accumulation, depuration and hepatic mixed-function oxidase enzyme induction in juvenile rainbow trout and lake whitefish exposed to dietary 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Aquat Toxicol* 37:201-220.

Guadagnolo, C.M., C.J. Brauner, and C.M. Wood. 2001. Chronic effects of silver exposure on ion levels, survival, and silver distribution within developing rainbow trout (*Oncorhynchus mykiss*) embryos. *Environ Toxicol Chem* 20(3):553-560.

Hansen DJ, Parrish PR, Lowe JL, Wilson AJ, Jr, Wilson PD. 1971. Chronic toxicity, uptake, and retention of Aroclor 1254 in two estuarine fishes. *Bull Environ Contam Toxicol* 6:113-119.

Hermanutz RO, Allen KN, Roush TH, Hedtke SF. 1992. Effects of elevated selenium concentrations on bluegills (*Lepomis macrochirus*) in outdoor experimnetal streams. *Environ Toxicol Chem* 11:217-224.

Holcombe, G.W., D.A. Benoit, E.N. Leonard, and J.M. McKim. 1976. Long-term effects of lead exposure on three generations of brook trout (*Salvelinus fontinalis*). *J Fish Res Bd Can* 33:1731-1741.

Lerner, D.T., B.T. Bjornsson, and S.D. McCormick, 2007. Effects of aqueous exposure to polychlorinated biphenyls (Aroclor 1254) on physiology and behavior of smolt development of Atlantic salmon, *Aquat. Toxicol.*, 81:329-336.

Mount, D.R., A.K. Barth, T.D. Garrison, K.A. Barten, and J.R. Hockett. 1994. Dietary and waterborne exposure of rainbow trout (*Oncorhynchus mykiss*) to copper, cadmium, lead and zinc using a live diet. *Environ Toxicol Chem* 13(12):2031-41.

Ogle RS, Knight AW. 1989. Effects of elevated foodborne selenium on growth and reproduction of the fathead minnow (*Pimephales promelas*). *Arch Environ Contam Toxicol* 18:795-803.

Parrish, P.R., S.C. Schimmel, D.J. Hansen, J.M. Patrick, Jr., and J. Forester. 1976. Chlordane: effects on several estuarine organisms. *J Toxicol Environ Health* 1:485-494.

Pierson KB. 1981. Effects of chronic zinc exposure on the growth, sexual maturity, reproduction, and bioaccumulation of the guppy, *Poecilia reticulata*. *Can J Fish Aquat Sci* 38:23-31

Schuytema, G.S., D.F. Krawczyk, W.L. Griffis, A.V. Nebeker, and M.L. Robideaux. 1990. Hexachlorobenzene uptake by fathead minnows and macroinvertebrates in recirculating sediment/water systems. *Arch Environ Contam Toxicol* 19:1-9.

Shubat, P.J. and L.R. Curtis. 1986. Ration and toxicant preexposure influence dieldrin accumulation by rainbow trout (*Salmo gairdneri*). *Environ Toxicol Chem* 5:69-77.

USEPA. 2014. Focused feasibility study report for the lower eight miles of the Lower Passaic River. The Louis Berger Group, Battelle, and HDR/HydroQual.

Webber HM, Haines TA. 2003. Mercury effects on predator avoidance behavior of a forage fish, golden shiner (*Notemigonus crysoleucas*). *Environ Toxicol Chem* 22(7):1556-1561

Windward Environmental LLC. 2019. Lower Passaic River Study Area Baseline Ecological Risk Assessment. Final. June

Zyadah, M.A. and T.E. Abdel-Baky. 2000. Toxicity and bioaccumulation of copper, zinc, and cadmium in some aquatic organisms. *Bull Environ Contam Toxicol* 64:740-747.

Table 7-11
TRVs for Fish Dietary Assessment

Constituent ^{a,b}	Units	Fish - Dietary TRVs		TRVs Developed Specifically for NBSA BERA	Source
		NOAEL	LOAEL		
PAHs					
Total PAH	mg/kg bw/day	6.2	18		Chinook salmon (anadromous); growth (Meador et al. 2006)
Inorganics					
Arsenic	mg/kg bw/day	0.52	1.9	X	Striped bass (marine/anad; growth; Blazer et al. 1997)
Cadmium	mg/kg bw/day	0.001 ^c	0.01		Rockfish (marine); growth (Kim et al. 2004); Kang et al. 2005)
Chromium	mg/kg bw/day	0.19	Not derived		Grey mullet (marine); growth (Walsh et al. 1994)
Copper	mg/kg bw/day	1	2	X	Rockfish (marine); growth (Kang et al. 2005)
Lead	mg/kg bw/day	12.6	Not derived		Rainbow trout (freshwater); growth (Mount et al. 1994)
Mercury/Methylmercury	mg/kg bw/day	0.00056 ^c	0.0056		10 species; growth, reproduction, mortality, and behavior (Windward Environmental LLC 2019)
Nickel	mg/kg bw/day	0.14 ^c	1.4		Indian carp (freshwater); growth (Javed 2013)
Selenium	mg/kg bw/day	0.011 ^c	0.11		7 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)
Silver	mg/kg bw/day	70	Not derived	X	Rainbow trout (freshwater; growth; Galvez and Wood 1999)
Zinc	mg/kg bw/day	19	38		Rainbow trout (freshwater); growth (Takeda and Shimma 1977)

Notes:

^a Risk for fish evaluated for metals and PAHs only.

^b The revised LPR FFS (USEPA 2014) for the lower eight miles of the Lower Passaic River did not include a fish dietary assessment.

^c NOAEL extrapolated from LOAEL using an uncertainty factor of 10.

Acronyms and Abbreviations:

FFS = focused feasibility study

LOAEL = lowest observed adverse effects level

LPR = Lower Passaic River

mg/kg bw/day = milligrams per kilogram body weight per day

NOAEL = no observed adverse effect level

PAH = polycyclic aromatic hydrocarbon

TRV = toxicity reference value

Table 7-11
TRVs for Fish Dietary Assessment

References

- Blazer, V.S., J.W. Fournie, and B.A. Weeks-Perkins. 1997. Macrophage aggregates: biomarker for immune function in fishes? In: Dwyer FJ, Doane TR, Hinman ML, eds, Environmental toxicology and risk assessment: Modeling and risk assessment. Vol 6. STP 1317. American Society for Testing and Materials, Philadelphia, PA, pp 360-375.
- Galvez, F. and C.M. Wood. 1999. Physiological effects of dietary silver sulfide exposure in rainbow trout. Environ Toxicol Chem 18(1):84-88.
- Hamilton, S.J., K.J. Buhl, N.L. Faerber, R.H. Wiedmeyer, and F.A. Bullard. 1990. Toxicity of organic selenium in the diet to chinook salmon. Environ Toxicol Chem 9:347-358.
- Hilton, J.W. and W.J. Bettger. 1988. Dietary vanadium toxicity in juvenile rainbow trout: a preliminary study. Aquat Toxicol 12:63-71.
- Javed, M. 2013. Chronic effects of nickel and cobalt on fish growth. Int J Agric Biol 15:575-579.
- Kang, J-C, S-G. Kim, and S-W. Jang. 2005. Growth and hematological changes of rockfish, *Sebastes schlegeli* (Hilgendorf) exposed to dietary Cu and Cd. J World Aquacult Soc 36(2):188-195.
- Kim, S-G, D-K. Park, S-W. Jang, J-S. Lee, S-S. Kim, and M-H. Chung. 2008. Effects of dietary benzo(a)pyrene on growth and hematological parameters in juvenile rockfish, *Sebastes schlegeli* (Hilgendorf). Bull Environ Contam Toxicol 81:470-474.
- Kim, S-G, J-W. Kim, and J-C. Kang. 2004. Effect of dietary cadmium on growth and haematological parameters of juvenile rockfish, *Sebastes schlegeli* (Hilgendorf). Aquacult Res 35:80-86.
- Matta, M.B., J. Linse, C. Cairncross, L. Francendese, and R.M. Kocan. 2001. Reproductive and transgenerational effects of methylmercury or Aroclor 1268 on *Fundulus heteroclitus*. Environ Toxicol Chem 20(2):327-335.
- Meador, J.P., F.C. Sommers, G.M. Ylitalo, and C.A. Sloan. 2006. Altered growth and related physiological responses in juvenile chinook salmon (*Oncorhynchus tshawytscha*) from dietary exposure to polycyclic aromatic hydrocarbons (PAHs). Can J Fish Aquat Sci 63:2364-2376.
- Mount, D.R., A.K. Barth, T.D. Garrison, K.A. Barten, and J.R. Hockett. 1994. Dietary and waterborne exposure of rainbow trout (*Oncorhynchus mykiss*) to copper, cadmium, lead and zinc using a live diet. Environ Toxicol Chem. 13(12):2031-41.
- Takeda H and Shimma Y. 1977. Effects of toxic amounts of dietary zinc on the growth and body components of rainbow trout at two levels of calcium. Bull Freshw Fish Res 27:103-109.
- Walsh, A.R., J. O'Halloran, and A.M. Gower. 1994. Some effects of elevated levels of chromium (III) in sediments to the mullet *Chelon labrosus* (R). Ecotox Environ Saf 27:168-176.
- Windward Environmental LLC. 2019. Lower Passaic River Study Area Baseline Ecological Risk Assessment. Final. June

Table 8-2
TRVs for Bird Dietary Assessment

Constituent	Units	Bird - Dietary TRVs						
		NBSA BERA TRV			TRVs Developed Specifically for NBSA BERA	LPR FFS TRV (USEPA 2014)		
		NOAEL	LOAEL	Source		NOAEL	LOAEL	Source
Dioxins/Furans								
2,3,7,8-TCDD	mg/kg bw/day	0.000014	1.4E-04	Ring-necked pheasant; mortality, growth, and reproduction (Nosek et al. 1992a)		0.0000028	0.000028	Ring-necked pheasant; mortality, growth, and reproduction (Nosek et al. 1992a, 1992b)
Total TEQ Bird	mg/kg bw/day	0.000014	0.00014	Ring-necked pheasant; mortality, growth, and reproduction (Nosek et al. 1992a)		0.0000028	0.000028	Ring-necked pheasant; mortality, growth, and reproduction (Nosek et al. 1992a, 1992b)
Polychlorinated Biphenyls								
Total PCBs	mg/kg bw/day	0.14 ^a	1.4	Ringed turtle-dove; reproduction (Peakall et al. 1972; Peakall and Peakall 1973)		0.4	0.5	Chicken; reproduction (Chapman 2003)
Total PCB TEQ Bird	mg/kg bw/day	0.000014	0.00014	Ring-necked pheasant; mortality, growth, and reproduction (Nosek et al. 1992a)		0.0000028	0.000028	Ring-necked pheasant; mortality, growth, and reproduction (Nosek et al. 1992a, 1992b)
Total PCB Dioxin TEQ Bird	mg/kg bw/day	0.000014	0.00014	Ring-necked pheasant; mortality, growth, and reproduction (Nosek et al. 1992a)		0.0000028	0.000028	Ring-necked pheasant; mortality, growth, and reproduction (Nosek et al. 1992a, 1992b)
Pesticides								
Total DDx (2,4 & 4,4)	mg/kg bw/day	0.025 ^a	0.25	10 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.009	0.027	Brown pelican; reproduction (Anderson et al. 1975)
Dieldrin	mg/kg bw/day	0.08	0.12	Quail; reproduction and mortality (DeWitt 1956)	X	0.054	0.18	Helmeted guinea fowl; survival (Wiese et al. 1969)
Total Chlordane	mg/kg bw/day	2 ^a	20	Bobwhite: mortality (Hill et al. 1976; Heath et al. 1972)	X	--	--	--
Hexachlorobenzene	mg/kg bw/day	1.1	5	Japanese quail; mortality (Vos et al. 1971)	X	--	--	--
Polycyclic Aromatic Hydrocarbons								
Total HMW PAH	mg/kg bw/day	--	--	--		0.048	0.48	Pigeon; reproduction (Hough et al. 1993)
Total LMW PAH	mg/kg bw/day	--	--	--		0.67	6.7	Red-winged blackbird; survival (Schafer et al. 1983)
Total PAH	mg/kg bw/day	40	Not derived	Mallard; growth (Patton and Dieter 1980)		--	--	--
Inorganics								
Arsenic	mg/kg bw/day	10	40	Mallard; reproduction (Stanley et al. 1994)	X	--	--	--
Cadmium	mg/kg bw/day	0.4 ^a	4	Japanese quail; growth (Richardson et al. 1974)		--	--	--
Chromium	mg/kg bw/day	10.5 ^a	105	Chicken; survival and growth (Chung et al. 1985)		--	--	--
Copper	mg/kg bw/day	1.9 ^a	19	Chicken; growth (Jensen and Maurice 1978)		2.3	4.7	Wild turkey; growth (Kashani et al. 1986)
Lead	mg/kg bw/day	5.5	28	Japanese quail; growth (Morgan et al. 1975)		0.19	1.9	Japanese quail; reproduction (Edens and Garlich 1983)
Mercury/Methylmercury	mg/kg bw/day	0.0096 ^a	0.096	6 species; survival, growth, and reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)		0.013	0.026	Mallard; reproduction (Heinz 1974, 1976, 1979)
Nickel	mg/kg bw/day	15	33	Chicken; growth (Weber and Reid 1968)		--	--	--
Selenium	mg/kg bw/day	0.42	0.82	Mallard; reproduction (Heinz et al. 1989)		--	--	--
Silver	mg/kg bw/day	9.86 ^a	98.6	Mallard; mortality (Van Vleet 1977 as cited in USEPA 2006)	X	--	--	--
Zinc	mg/kg bw/day	82	124	Chicken; growth (Roberson and Schaible 1960)		--	--	--

Table 8-2
TRVs for Bird Dietary Assessment

Notes:
^a NOAEL extrapolated from LOAEL using an uncertainty factor of 10.

Acronyms and Abbreviations:

BERA = baseline ecological risk assessment	NOAEL = no observed adverse effect level
COPEC = constituent of potential ecological concern	PAH = polycyclic aromatic hydrocarbon
FFS = focused feasibility study	PCB = polychlorinated biphenyl
HMW = high molecular weight	SSD = species sensitivity distribution
LMW = low molecular weight	TCDD = tetrachlorodibenzo-p-dioxin
LPR = Lower Passaic River	TEQ = toxic equivalent
LPRSA = Lower Passaic River Study Area	Total DDx = sum of dichlorodiphenyldichloroethylene, dichlorodiphenyldichloroethane, and dichlorodiphenyltrichloroethane
LOAEL = lowest observed adverse effects level	TRV = toxicity reference value
mg/kg bw/day = milligrams per kilogram of body weight per day	USEPA = U.S. Environmental Protection Agency
NBSA = Newark Bay Study Area	-- = not available

References

Anderson, D.W., R.W. Risebrough, L.A. Woods Jr., L.R. DeWeese, and W.G. Edgecomb. 1975. Brown pelicans: improved reproduction off the southern California coast. Science 190:806-808.

Chapman, J. 2003. Memorandum dated March 6, 2003: Toxicity reference values (TRVs) for mammals and birds based on selected Aroclors. Ecologist, US Environmental Protection Agency Region 5, Chicago, IL.

Chung, K.H., Y.O. Sukgrand, and M.H. Kang. 1985. The toxicity of chromium and its interaction with manganese and molybdenum in chicks. Kor J Anim Sci 27(6):391-395.

Davison, K.L. and J.L. Sell. 1974. DDT thins shells of eggs from mallard ducks maintained on ad libitum or controlled-feeding regimens. Arch Environ Contam Toxicol 2(3):222-232.

Edens, F.W. and J.D. Garlich. 1983. Lead-induced egg production decrease in leghorn and Japanese quail hens. Poult Sci 62:1757-1763.

Heath, R., G., J.W. Spann, E.F. Hill, and J.F. Kreitzer. 1972. Comparative Dietary Toxicity of Pesticides to Birds. Special Scientific Report. Wildl.No.152. 57 pgs.

Heinz, G.H. 1974. Effects of low dietary levels of methyl mercury on mallard reproduction. Bull Environ Contam Toxicol 11(4):386-392.

Heinz, G.H. 1976. Methylmercury: second-generation reproductive and behavioral effects on mallard ducks. J Wildl Manage 40(4):710-715

Heinz, G.H. 1979. Methylmercury: reproductive and behavioral effects on three generations of mallard ducks. J Wildl Manage 43(2):394-401.

Heinz, G.H., D.J. Hoffman, and L.G. Gold. 1989. Impaired reproduction of mallards fed an organic form of selenium. J Wildl Manage 53(2):418-428.

Hill, E.F., R.G. Heath, and J.D. Williams. 1976. Effect of Dieldrin and Aroclor 1242 on Japanese Quail Eggshell Thickness. Bull. Environ. Contam. Toxicol. 16(4): 445-453. Ref ID: 995 .

Hough, J.L., M.B. Baird, G.T. Sfeir, C.S. Pacini, D. Darrow, and C. Wheelock. 1993. Benzo(a)pyrene enhances atherosclerosis in white carneau and show racer pigeons. Arterioscler Thromb 13:1721-1727.

Jensen, L.S. and D.V. Maurice. 1978. Effect of methionine on copper-induced growth depression and gizzard erosion. Poult Sci 57:1530-1532.

Kashani, A.B., H. Samie, R.J. Emerick, and C.W. Carlson. 1986. Effect of copper with three levels of sulfur containing amino acids in diets for turkeys. Poult Sci 65:1754-1759.

Nosek, J.A., J.R. Sullivan, T.E. Amundson, S.R. Craven, L.M. Miller, A.G. Fitzpatrick, M.E. Cook, and R.E. Peterson, 1992b. Embryotoxicity of 2,3,7,8 tetrachlorodibenzo-p-dioxin in ring-necked pheasants; Environ Contam Toxicol 12:1215-1222.

Nosek, J.A., S.R. Craven, J.R. Sullivan, S.S. Hurley, and R.E. Peterson. 1992a. Toxicity and reproductive effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin in ring-necked pheasant hens. J Toxicol Environ Health 35:187-198.

Patton, J.F. and M.P. Dieter. 1980. Effects of petroleum hydrocarbons on hepatic function in the duck. Comp Biochem Physiol 65C:33-36.

Peakall, D.B. and M.L. Peakall. 1973. Effect of a polychlorinated biphenyl on the reproduction of artificially and naturally incubated dove eggs. J Appl Ecol 10:863-868.

Peakall, D.B., J.L. Lincer, and S.E. Bloom. 1972. Embryonic mortality and chromosomal alterations caused by Aroclor 1254 in ring doves. Environ Health Perspect 1:103-104

Richardson, M.E., M.R.S. Fox, and B.E. Bry, Jr. 1974. Pathological changes produced in Japanese quail by ingestion of cadmium. J Nutr 104:323-338.

Roberson RH, Schaible PJ. 1960. The tolerance of growing chicks for high levels of different forms of zinc. Poult Sci 39:893-895.

Schafer, E.W., Jr, W.A. Bowles, Jr, and J. Hurlbut. 1983. The acute oral toxicity, repellency, and hazard potential of 998 chemicals to one or more species of wild and domestic birds. Arch Environ Contam Toxicol 12:355-382.

USEPA. 2006. Ecological soil screening levels for silver. Interim final. OSWER Directive 9285.7-77. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington. Available from: <https://rais.ornl.gov/documents/>

USEPA. 2014. Focused feasibility study report for the lower eight miles of the Lower Passaic River. The Louis Berger Group, Battelle, and HDR/HydroQual

Van Vleet, John F. 1977. Protection by Various Nutritional Supplements Against Lesions of Selenium-vitamin E Deficiency Induced in Ducklings Fed Tellurium or Silver. Am. J. Vet. Res. 38(9): 1393-8.

Vos, J.G., H.L. Van der Maas, A. Musch, and E. Pam. 1971. Toxicity of hexachlorobenzene in Japanese quail with spcial reference to porphyria, liver damage, reproduction, and tissue residues. Toxicology and Applied Pharmacology 18: 944-957.

Weber, C.W. and B.L. Reid. 1968. Nickel toxicity in growing chicks. J Anim Sci 95:612-616.

Windward Environmental LLC. 2019. Lower Passaic River Study Area Baseline Ecological Risk Assessment. Final. June

Wiese, I.H., N.C.J. Basson, J.H. Van Der Vyver, and J.H. Van Der Merwe. 1969. Toxicology and Dynamics of Dieldrin in the Crowned Guinea-fowl, Numida Meleagris (L). Phytophylactica 1(3-4): 161-175. Ref ID: 1158.

Table 8-5
TRVs for Bird Egg Tissue Assessment

Constituent	Units	Bird - Egg TRVs					
		NBSA BERA TRV			LPR FFS TRV (USEPA 2014)		
		NOAEL	LOAEL	Source	NOAEL	LOAEL	Source
Dioxins/Furans							
2,3,7,8-TCDD	mg/kg, ww	0.000025 ^a	0.00025	5 species; reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)	0.000059	0.00015	Various species; reproduction (USEPA 2003)
Total Dioxin/Furan TEQ Bird	mg/kg, ww	0.000025 ^a	0.00025	5 species; reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)	0.000059	0.00015	Various species; reproduction (USEPA 2003)
Polychlorinated Biphenyls							
Total PCBs	mg/kg, ww	1.6 ^a	16	Ringed turtle dove; reproduction (Peakall et al. 1972); Peakall and Peakall 1973)	0.7	1.3	Chicken; reproduction (Chapman 2003)
Total PCB TEQ Bird	mg/kg, ww	0.000025 ^a	0.00025	5 species; reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)	0.000059	0.00015	Various species; reproduction (USEPA 2003)
Total Dioxin/Furan/PCB TEQ Bird	mg/kg, ww	0.000025 ^a	0.00025	5 species; reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)	0.000059	0.00015	Various species; reproduction (USEPA 2003)
Pesticides							
Total DDx (2,4 & 4,4)	mg/kg, ww	0.41 ^a	4.1	7 species; reproduction. SSD-derived 5th percentile value. (Windward Environmental LLC 2019)	0.5	3	Brown pelican; reproduction (Blus 1984)
Dieldrin	mg/kg, ww	0.3 ^a	3	Pheasant; reproduction (Genelly and Rudd 1956)	0.2	8.1	Barn owl; reproduction (Mendenhall et al. 1983)
Inorganics							
Mercury/Methylmercury	mg/kg, ww	0.18 ^a	1.8	Mallard; reproduction (Heinz 1979, Heinz and Hoffman 2003, Heinz 1974, 1976)	0.011	0.11	Carolina wren; 'Reproduction (Jackson 2011)

Table 8-5
TRVs for Bird Egg Tissue Assessment

Notes

^a NOAEL extrapolated from LOAEL using an uncertainty factor of 10.

No bird egg TRVs were developed specifically for the NBSA BERA (Windward Environmental LLC 2019). All TRVs were selected from the LPRSA BERA or the LPR FFS (USEPA 2003).

Acronyms and Abbreviations:

BERA = baseline ecological risk assessment

FFS = focused feasibility study

LOAEL = lowest observed adverse effects level

LPR = Lower Passaic River

mg/kg, ww = milligrams per kilogram, wet weight

NBSA = Newark Bay Study Area

NOAEL = no observed adverse effect level

PCB = polychlorinated biphenyl

SSD = species sensitivity distribution

TCDD = tetrachlorodibenzo-p-dioxin

TEQ = toxic equivalent

Total DDx = sum of dichlorodiphenyldichloroethylene, dichlorodiphenyldichloroethane, and dichlorodiphenyltrichloroethane

TRV = toxicity reference value

USEPA = U.S. Environmental Protection Agency

-- = not available

References:

- Blus, L.J. 1984. DDE in birds' eggs: comparison of two methods for estimating critical levels. *Wilson Bull* 96(2):268-276.
- Chapman, J. 2003. Memorandum dated March 6, 2003: Toxicity reference values (TRVs) for mammals and birds based on selected Aroclors. Ecologist, US Environmental Protection Agency Region 5, Chicago, IL.
- Genelly, R.E. and R.L. Rudd. 1956. Effects of DDT, toxaphene, and dieldrin on pheasant reproduction. *Auk* 73:529-539.
- Heinz, G.H. 1974. Effects of low dietary levels of methyl mercury on mallard reproduction. *Bull Environ Contam Toxicol* 11(4):386-392.
- Heinz, G.H. 1976. Methylmercury: second-year feeding effects on mallard reproduction and duckling behavior. *J Wildl Manag* 40(1):82-90.
- Heinz, G.H. 1979. Methylmercury: reproductive and behavioral effects on three generations of mallard ducks. *J Wildl Manag* 43(2):394-401.
- Heinz, G.H. and D.J. Hoffman. 2003. Embryotoxic thresholds of mercury: estimates from individual mallard eggs. *Arch Environ Contam Toxicol* 44:257-264.
- Jackson, A.K., D.C. Evers, M.A. Ettersson, A.M. Condon, S.B. Folsom, J. Detweiler, J. Schmerfeld, D.A. Cristol. 2011. Mercury exposure affects the reproductive success of a free-living terrestrial songbird, the Carolina wren (*Thryothorus ludovicianus*). *Auk* 128:759-769.
- Mendenhall, V.M., E.E. Klaas, and M.A.R. McLane. 1983. Breeding success of barn owls (*Tyto alba*) fed low levels of DDE and dieldrin. *Arch Environ Contam Toxicol* 12:235-240.
- Peakall, D.B. and M.L. Peakall. 1973. Effect of a polychlorinated biphenyl on the reproduction of artificially and naturally incubated dove eggs. *J App Ecol* 10:863-868.
- Peakall, D.B., J.L. Lincer, and S.E. Bloom. 1972. Embryonic mortality and chromosomal alterations caused by Aroclor 1254 in ring doves. *Environ Health Perspect* 1:103-104.
- USEPA. 2003. Analyses of laboratory and field studies of reproductive toxicity in birds exposed to dioxin-like compounds for use in ecological risk assessment. EPA/600/R-03/114F. National Center for Environmental Assessment, US Environmental Protection Agency, Cincinnati, OH.
- USEPA. 2014. Focused feasibility study report for the lower eight miles of the Lower Passaic River. The Louis Berger Group, Battelle, and HDR/HydroQual.
- Windward Environmental LLC. 2019. Lower Passaic River Study Area Baseline Ecological Risk Assessment. Final. June

Table 9-3
TRVs for Mammal Dietary Assessment

Constituent	Units	Mammal - Dietary TRVs						
		NBSA BERA TRV			TRVs Developed Specifically for NBSA BERA	LPR FFS TRV (USEPA 2014)		
		NOAEL	LOAEL	Source		NOAEL	LOAEL	Source
Dioxins/Furans								
2,3,7,8-TCDD	mg/kg bw/day	0.0000026	0.0000088	Mink; reproduction (Hochstein et al. 2001)		0.00000008	0.0000022	Mink; reproduction (Tillitt et al. 1996)
Total Dioxin/Furan TEQ Mammal	mg/kg bw/day	0.0000026	0.0000088	Mink; reproduction (Hochstein et al. 2001)		0.00000008	0.0000022	Mink; reproduction (Tillitt et al. 1996)
Polychlorinated Biphenyls								
Total PCBs	mg/kg bw/day	0.08	0.10	Mink; reproduction (Chapman 2003)		0.069	0.082	Mink; reproduction (Chapman 2003)
Total PCB TEQ Mammal	mg/kg bw/day	0.0000026	0.0000088	Mink; reproduction (Hochstein et al. 2001)		0.00000008	0.0000022	Mink; reproduction (Tillitt et al. 1996)
Total Dioxin/Furan/PCB TEQ Mammal	mg/kg bw/day	0.0000026	0.0000088	Mink; reproduction (Hochstein et al. 2001)		0.00000008	0.0000022	Mink; reproduction (Tillitt et al. 1996)
Pesticides								
Total DDX (2,4 & 4,4)	mg/kg bw/day	0.13 ^a	1.30	Mouse; reproduction (Ware and Good 1967)	X	0.8	4	Rat; reproduction (Fitzhugh 1948)
Dieldrin	mg/kg bw/day	0.02	0.03	Rat; reproduction (Harr et al. 1970)		0.015	0.03	Rat; reproduction (Harr et al. 1970)
Chlordane	mg/kg bw/day	2.1 ^a	21	Rat; reproduction and growth (Narotsky and Kavlok 1995)	X	--	--	--
Hexachlorobenzene	mg/kg bw/day	0.013 ^a	0.13	Mink; reproduction (Bleavins et al. 1984)	X	--	--	--
Polycyclic Aromatic Hydrocarbons								
Total HMW PAH	mg/kg bw/day	--	--	--		0.62	3.1	Mouse; survival (Culp et al. 1998)
Total LMW PAH	mg/kg bw/day	--	--	--		50	150	Rat; growth (Navarro et al. 1991)
Total PAH	mg/kg bw/day	0.615	3.07	Mouse; survival (Culp et al 1998 as cited in USEPA 2007c)	X	--	--	--
Inorganics								
Arsenic	mg/kg bw/day	2.6	5.4	Rat; growth (Byron et al. 1967)		--	--	--
Cadmium	mg/kg bw/day	3.5	13	Rat; growth (Machemer and Lorke 1981)		--	--	--
Chromium	mg/kg bw/day	0.96 ^a	9.62	Mouse; reproduction (Zahid et al. 1990; cited in USEPA 2008)	X	--	--	--
Copper	mg/kg bw/day	18	26	Mink; reproduction (Aulerich et al. 1982)		3.4	6.8	Mink; reproduction (Aulerich et al. 1982; cited in USEPA 2007a)
Lead	mg/kg bw/day	11	90	Rat; growth (Azar et al. 1973)		0.71	7	Rat; reproduction (Grant et al. 1980; cited in USEPA 2005b)
Mercury/Methylmercury	mg/kg bw/day	0.16	0.25	Mink; growth/survival (Wobeser et al. 1976)		0.016	0.027	Mink; growth and survival (Wobeser et al. 1976; cited in USEPA 1995)
Nickel	mg/kg bw/day	40	80	Rat; reproduction (Ambrose et al. 1976)		--	--	--
Selenium	mg/kg bw/day	0.016 ^a	0.16	Rat; growth (Behne et al. 1992)		--	--	--
Silver	mg/kg bw/day	18.8 ^a	188	Rat; reproduction (Shavlovski et al. 1995; cited in USEPA 2006)	X	--	--	--
Zinc	mg/kg bw/day	160	320	Rat; reproduction (Schlicker and Cox 1968)		--	--	--

Table 9-3
TRVs for Mammal Dietary Assessment

Notes:

^a NOAEL extrapolated from LOAEL using an uncertainty factor of 10.

Acronyms and Abbreviations:

BERA = baseline ecological risk assessment	NOAEL = no observed adverse effect level
COPEC = constituent of potential ecological concern	PAH = polycyclic aromatic hydrocarbon
FFS = focused feasibility study	PCB = polychlorinated biphenyl
HMW = high molecular weight	SSD = species sensitivity distribution
LMW = low molecular weight	TCDD = tetrachlorodibenzo-p-dioxin
LPR = Lower Passaic River	TEQ = toxic equivalent
LPRSA = Lower Passaic River Study Area	Total DDx = sum of dichlorodiphenyldichloroethylene (DDE), dichlorodiphenyldichloroethane (DDD), and dichlorodiphenyltrichloroethane (DDT)
LOAEL = lowest observed adverse effects level	TRV = toxicity reference value
mg/kg bw/day = milligrams per kilogram of body weight per day	USEPA = U.S. Environmental Protection Agency
NBSA = Newark Bay Study Area	-- = not available

References:

Ambrose AM, Larson PS, Borzelleca JF, Hennigar Jr GR. 1976. Long term toxicologic assessment of nickel in rats and dogs. J Food Sci Technol 13(4):181-187.

Aulerich RJ, Ringer RK, Bleavins MR, Napolitano A. 1982. Effects of supplemental dietary copper on growth, reproductive performance and kit survival of standard dark mink and the acute toxicity of copper to mink. J Anim Sci 55(2):337-343.

Aulerich, R.J., R.K. Ringer, M.R. Bleavins, and A. Napolitano. 1982. Effects of supplemental dietary copper on growth, reproductive performance and kit survival of standard dark mink and the acute toxicity of copper to mink. J Anim Sci 55(2):337-343.

Azar A, Trochimowicz HJ, Maxfield ME. 1973. Review of lead studies in animals carried out at Haskell Laboratory-two-year feeding study and response to hemorrhage study. In: Barth D, ed. Environmental Health Aspects of Lead, International Symposium, Amsterdam, the Netherlands. Commission of European Communities. pp 199-210.

Behne D, Kyriakopoulos A, Gessner H, Walzog B, Meinhold H. 1992. Type I iodothyronine deiodinase activity after high selenium intake, and relations between selenium and iodine metabolism in rats. J Nutr 122(7):1542-1546.

Bleavins, M.R., R.J. Aulerich, and R.K. Ringer. 1984. Effects of chronic dietary hexachlorobenzene exposure on the reproductive performance and survivability of mink and European ferrets. Arch Environ Contam Toxicol 13:357-365.

Byron, W.R., G.W. Bierbower, J.B. Brouwer, W.H. Hansen. 1967. Pathologic changes in rats and dogs from a two-year feeding of sodium arsenite or sodium arsenate. Toxicology and Applied Pharmacology 10: 132-147.

Chapman, J. 2003. Memorandum dated March 6, 2003: Toxicity reference values (TRVs) for mammals and birds based on selected Aroclors. Ecologist, US Environmental Protection Agency Region 5, Chicago, IL.

Culp, S.J., D.W. Gaylor, W.G. Sheldon, L.S. Goldstein, and F.A. Beland. 1998. A comparison of the tumors induced by coal tar and benzo[a]pyrene in a 2-year bioassay. Carcinogenesis 19(1):117-124.

Fitzhugh, O.G. 1948. Use of DDT insecticides on food products. Ind and Eng Chemistry 40(4):704-705.

Grant, L. D., C.A. Kimmel, G.L. West, C.M. Martinez-Vargas, and J.L. Howard. 1980. Chronic Low Level Lead Toxicity in the Rat. 2. Effects on Postnatal Physical and Behavioral Development. Toxicol Appl Pharmacol 56(1):42-58.

Harr, J.R., R.R. Claeys, J.F. Bone, and T.W. McCorcle. 1970. Dieldrin toxidosis: rat reproduction. Am J Vet Res 31(1):181-189.

Hext PM, Pinto PJ, Rimmel BA. 1999. Subchronic feeding study of antimony trioxide in rats. J Appl Toxicol 19:205-209.

Hochstein, J.R., J.A. Render, S.J. Bursian, and R.J. Aulerich. 2001. Chronic toxicity of dietary 2,3,7,8-tetrachlorodibenzo-p-dioxin to mink. Vet Hum Toxicol 43(3):134-139.

Machemer L, Lorke D. 1981. Embryotoxic effect of cadmium on rats upon oral administration. Toxicol Appl Pharmacol 58:438-443.

Narotskv, M.G. and R.J. Kavlock. 1995. A multidisciplinary approach to toxicological screening: II. Developmental toxicity. J Toxicol Environ Health 1995;45(2):145-71.

Navarro, HA; Price, CJ; Marr, MC; Myers, CB; Heindel, JJ; Schwetz, BA. 1991. Developmental toxicity evaluation of naphthalene (CAS NO. 91-20-3) administered by gavage to Sprague-Dawley (CD) rats on gestation days 6 through 15. Chemistry and Life Sciences, Research Triangle Institute (RTI). Research Triangle Park, NC. RTI Master Protocol Number: RTI-376. December 5.

Schlicker, S.A. and D.H. Cox. 1968. Maternal dietary zinc, and development and zinc, iron, and copper content of the rat fetus. J Nub 95:287-294.

Shavlovski, M.M., N.A. Chebotar, L.A. Konopistseva, E.T. Zakharova, A.M. Kachourin, V.B. Vassiliev, and V.S. Gaitskhoki. 1995. Embryotoxicity of Silver Ions is Diminished by Ceruloplasmin-further Evidence for its Role in the Transport of Copper. Biometals 8(2):122-8.

Tillitt, D.E., R.W. Gale, J.C. Meadows, J.L. Zajicek, P.H. Peterman, S.N. Heaton, P.D. Jones, S.J. Bursian, T.J. Kubiak, J.P. Giesy, and R.J. Aulerich. 1996. Dietary exposure of mink to carp from Saginaw Bay. 3. Characterization of dietary exposure to planar halogenated hydrocarbons, dioxin equivalents, and biomagnification. Environ Sci Technol 30:283-291.

USEPA. 1995. Great Lakes Water Quality Initiative criteria documents for the protection of wildlife: DDT, mercury, 2,3,7,8-TCDD, PCBs. EPA-820-B-95-008. Office of Water, US Environmental Protection Agency, Washington, DC.

USEPA. 2005b. Ecological soil screening levels for lead. Interim final. OSWER Directive 9285.7-70. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC. Available from: https://rais.ornl.gov/documents/eco-ssl_lead.pdf.

USEPA. 2006. Ecological soil screening levels for silver. Interim final. OSWER Directive 9285.7-77. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington. Available from: https://rais.ornl.gov/documents/eco-ssl_silver.pdf

USEPA. 2007a. Ecological soil screening levels for copper. Interim final. Revised February 2007. OSWER Directive 9285.7-68. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC. Available from: https://rais.ornl.gov/documents/eco-ssl_copper.pdf

USEPA. 2007c. Ecological soil screening levels for Polycyclic Aromatic Hydrocarbons (PAHs). Interim final. OSWER Directive 9285.7-78. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington. Available from: https://rais.ornl.gov/documents/eco-ssl_pah.pdf.

USEPA. 2008. Ecological soil screening levels for chromium. Interim final. OSWER Directive 9285.7- 66. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC. Available from: https://rais.ornl.gov/documents/eco-ssl_chromium.pdf

USEPA. 2014. Focused feasibility study report for the lower eight miles of the Lower Passaic River. The Louis Berger Group, Battelle, and HDR/HydroQua

Ware, G.W. and E.E. Good. 1967. Effects of insecticides on reproduction in the laboratory mouse. II. Mirex, telodrin, and DDT. Toxicol Appl Pharmacol 10:54-61.

Wobeser, G., N.D. Nielson, and B. Schiefer. 1976. Mercury and Mink II: Experimental methyl mercury intoxication. Can J Comp Med 40:34-45.

Zahid, Z.R., Z.S. Al-Hakkak, A.H.H. Kadhim, E.A. Elias, and I.S. Al-Jumaily. 1990. Comparative effects of trivalent and hexavalent chromium on spermatogenesis of the mouse. Toxicol Environ Chem 25(2-3):131-136.

Table 11-1a
Weight of Evidence Risk Summary for NBSA BERA

Receptor	Preliminary COC ^a	Primary LOE	Summary of LOE	Risk Conclusions	Risk Driver ^b
Assessment Endpoint: Survival, growth, and/or reproduction of benthic invertebrates					
Benthic Invertebrate Communities	see Table 11-1b for Sediment Quality Triad risk conclusions			Localized Risk per SQT	
Polychaete Worm	2,3,7,8-TCDD, Total PCBs, Arsenic	SQT	<ul style="list-style-type: none"> Localized risk to benthic communities (see Table 11-1b for conclusions of SQT) No effect on growth/survival in site-specific toxicity tests (Iannuzzi et al. 2008) 	Localized Risk per SQT	
Softshell Clam	2,3,7,8-TCDD, Total PCBs	Tissue HQs	<ul style="list-style-type: none"> LPR FFS HQs ≥ 1 LPR FFS TRVs based on bivalve data Site-specific study on eastern oyster indicates potential bivalve reproductive impairment (Wintermyer and Cooper 2003) 	Possible	Yes
Softshell Clam	Arsenic, Lead	Tissue HQs	<ul style="list-style-type: none"> High magnitude of the HQs ≥ 1 High uncertainty with invertebrate tissue HQ approach 	Possible	Yes
Softshell Clam	Mercury	Tissue HQs	<ul style="list-style-type: none"> NBSA and LPR FFS HQs ≥ 1 in all scenarios Low magnitude of the HQs ≥ 1 Invertebrate tissue HQ approach for mercury/methylmercury less uncertain 	Possible	Yes
Softshell Clam	Total HMW PAHs, Cadmium, Chromium, Copper, Nickel, Selenium, Silver, Zinc	Tissue HQs	<ul style="list-style-type: none"> Low magnitude of the HQs ≥ 1 High uncertainty with invertebrate tissue HQ approach Selenium and silver present at concentrations near or below background 	Unlikely	No
Blue Crab	2,3,7,8-TCDD, Total PCBs	Tissue HQs	<ul style="list-style-type: none"> LPR FFS HQs ≥ 1 High magnitude of the HQs ≥ 1 LPR FFS TRVs based on bivalve data and considered uncertain for crabs 	Possible	Yes
Blue Crab	Total DDx	Tissue HQs	<ul style="list-style-type: none"> High magnitude of the HQs ≥ 1 based on alternate NBSA TRVs High uncertainty with invertebrate tissue HQ approach 	Possible	Yes
Blue Crab	Mercury, Methylmercury	Tissue HQs	<ul style="list-style-type: none"> NBSA and LPR FFS HQs ≥ 1 in all scenarios Low magnitude of the HQs ≥ 1 Invertebrate tissue HQ approach for mercury/methylmercury less uncertain 	Possible	Yes
Blue Crab	Dieldrin, Arsenic, Cadmium, Copper, Mercury, Methylmercury, Selenium, Silver	Tissue HQs	<ul style="list-style-type: none"> Low magnitude of the HQs ≥ 1 High uncertainty with invertebrate tissue HQ approach Selenium and silver present at concentrations near or below background 	Unlikely	No

Table 11-1a
Weight of Evidence Risk Summary for NBSA BERA

Receptor	Preliminary COC ^a	Primary LOE	Summary of LOE	Risk Conclusions	Risk Driver ^b
Assessment Endpoint: Survival, growth, and/or reproduction of fish					
Mummichog/Killifish	Dioxins/Furans and TEQ (2,3,7,8-TCDD, Total Dioxin/Furan TEQ Fish, PCB TEQ Fish, Dioxin/Furan/PCB TEQ Fish)	Tissue HQs	<ul style="list-style-type: none"> • LPR FFS WB tissue HQs > 10 • Alternate NBSA WB tissue HQs > 1 • NBSA WB tissue HQs < 1 • Egg HQs < 1 • Site-specific pathology data did not show abnormalities • Site-specific reproductive study indicates endocrine impariments relative to populations that are not urban-adapted (Bugel et al. 2010, 2011, 2014) 	Possible	Yes
Mummichog/Killifish	Total PCBs	Tissue HQs	<ul style="list-style-type: none"> • Tissue LOAEL HQs < 1 for most scenarios • Egg HQs < 1 • Low magnitude of the HQs ≥ 1 • Site-specific pathology data did not show abnormalities 	Unlikely	No
Mummichog/Killifish	Total DDx (2,4 & 4,4)	Tissue HQs	<ul style="list-style-type: none"> • LOAEL HQs < 1 in most scenarios • Low magnitude of HQs ≥ 1 in Southwest Zone • Site-specific pathology data did not show abnormalities 	Unlikely	No
Mummichog/Killifish	Cadmium	Tissue HQs	<ul style="list-style-type: none"> • Cadmium not detected in fish tissue • Low magnitude of dietary HQs ≥ 1 • Diet assumptions overestimate dietary exposure • Site-specific pathology data did not show abnormalities 	Unlikely	No
Mummichog/Killifish	Copper	Background	<ul style="list-style-type: none"> • Tissue copper at/within background • Dietary HQs < 1 • Low magnitude of tissue HQs ≥ 1 • Diet assumptions overestimate dietary exposure • Site-specific pathology data did not show abnormalities 	Unlikely	No
Mummichog/Killifish	Mercury, Methylmercury	Tissue HQs	<ul style="list-style-type: none"> • WB Tissue HQs < 1 • Dietary HQs < 1 in most scenarios • Low magnitude of dietary HQs ≥ 1 in Southwest Zone • Egg LOAEL HQs < 1 • Site-specific pathology data did not show abnormalities 	Unlikely	No
Mummichog/Killifish	Silver	Background	<ul style="list-style-type: none"> • Tissue LOAEL HQs < 1 in most scenarios • Silver < background in sediment • Low magnitude of tissue HQs ≥ 1 in North Zone • Dietary LOAEL HQs < 1 • Low confidence in both tissue and dietary TRVs • Site-specific pathology data did not show abnormalities 	Unlikely	No
American Eel	Dioxins/Furans and TEQ (2,3,7,8-TCDD, Total Dioxin/Furan TEQ Fish, Dioxin/Furan/PCB TEQ Fish)	Tissue HQs	<ul style="list-style-type: none"> • LPR FFS WB tissue HQs approaching 10 • NBSA WB tissue HQs < 1 • Site-specific pathology data did not show abnormalities • Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Possible	Yes

Table 11-1a
Weight of Evidence Risk Summary for NBSA BERA

Receptor	Preliminary COC ^a	Primary LOE	Summary of LOE	Risk Conclusions	Risk Driver ^b
American Eel	Total PCBs	Tissue HQs	<ul style="list-style-type: none"> NBSA WB tissue HQs < 1 Low magnitude of the WB tissue HQs ≥ 1 based on LPR FFS TRVs Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
American Eel	Total DDx (2,4 & 4,4)	Tissue HQs	<ul style="list-style-type: none"> NBSA WB tissue HQs < 1 Low magnitude of the WB tissue HQs ≥ 1 based on LPR FFS TRVs in North Zone Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
American Eel	Cadmium	Tissue HQs	<ul style="list-style-type: none"> Cadmium not detected in fish tissue Low magnitude of dietary HQs ≥ 1 Diet assumptions overestimate dietary exposure Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
American Eel	Copper	Dietary HQs	<ul style="list-style-type: none"> Dietary HQs < 1 for all scenarios Low magnitude of tissue HQs ≥ 1 in Southwest Zone Diet assumptions overestimate dietary exposure Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
American Eel	Mercury/Methylmercury	Tissue HQs	<ul style="list-style-type: none"> Tissue LOAEL HQs < 1 in most scenarios Dietary LOAEL HQs < 1 in most scenarios Low magnitude of LOAEL HQs ≥ 1 Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
American Eel	Selenium	Background	<ul style="list-style-type: none"> Selenium < background in sediment Tissue LOAEL HQs < 1 in most scenarios Low magnitude of tissue HQs ≥ 1 in Southeast Zone Dietary HQs < 1 Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No

Table 11-1a
Weight of Evidence Risk Summary for NBSA BERA

Receptor	Preliminary COC ^a	Primary LOE	Summary of LOE	Risk Conclusions	Risk Driver ^b
Summer/Winter Flounder	Dioxins/Furans and TEQ (2,3,7,8-TCDD, Total Dioxin/Furan TEQ Fish, Dioxin/Furan/PCB TEQ Fish)	Tissue HQs	<ul style="list-style-type: none"> LPR FFS WB tissue HQs approaching 10 NBSA WB tissue HQs < 1 Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Possible	Yes
Summer/Winter Flounder	Total PCBs	Tissue HQs	<ul style="list-style-type: none"> LOAEL HQs < 1 for most scenarios Low magnitude of the HQs ≥ 1 in the North Zone Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
Summer/Winter Flounder	Cadmium	Tissue HQs	<ul style="list-style-type: none"> Cadmium not detected in fish tissue Low magnitude of dietary HQs ≥ 1 Dietary HQs in Southwest Zone calculated using clam MDC Diet assumptions overestimate dietary exposure Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
Summer/Winter Flounder	Copper	Dietary HQs	<ul style="list-style-type: none"> Dietary HQs < 1 WB tissue HQs < 1 in most scenarios Low magnitude of tissue HQs ≥ 1 in North Zone Diet assumptions overestimate dietary exposure Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
Summer/Winter Flounder	Mercury/Methylmercury	Tissue HQs	<ul style="list-style-type: none"> Tissue LOAEL HQs < 1 Low magnitude of dietary HQs ≥ 1 Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
White Perch	Dioxins/Furans and TEQ (2,3,7,8-TCDD, Total Dioxin/Furan TEQ Fish, Dioxin/Furan/PCB TEQ Fish)	Tissue HQs	<ul style="list-style-type: none"> WB tissue LOAEL HQs ≥ 1 for most scenarios WB tissue LPR FFS LOAEL HQs > 10 for all scenarios Liver LOAEL HQs < 1 Liver HQs superseded by WB tissue HQs Site-specific pathology data did not show abnormalities Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Possible	Yes

Table 11-1a
Weight of Evidence Risk Summary for NBSA BERA

Receptor	Preliminary COC ^a	Primary LOE	Summary of LOE	Risk Conclusions	Risk Driver ^b
White Perch	Total PCB TEQ, Total PCBs	Tissue HQs	<ul style="list-style-type: none"> • WB tissue LOAEL HQs < 1 in most scenarios • Low magnitude of the WB tissue HQs ≥ 1 • Liver LOAEL HQs < 1 • Liver HQs superseded by WB tissue HQs • Site-specific pathology data did not show abnormalities • Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
White Perch	Total DDx (2,4 & 4,4)	Tissue HQs	<ul style="list-style-type: none"> • WB tissue LOAEL HQs > 1 in most cases • Low magnitude of the tissue LOAEL HQ ≥ 1 • Site-specific pathology data did not show abnormalities • Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Possible	Yes
White Perch	Arsenic, Zinc	Tissue HQs	<ul style="list-style-type: none"> • Tissue and dietary HQs < 1 • Low magnitude of the liver HQs ≥ 1 (arsenic) or based on unbounded NOAEL (zinc) • Liver HQs superseded by WB tissue HQs • Site-specific pathology data did not show abnormalities • Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
White Perch	Cadmium	Tissue HQs	<ul style="list-style-type: none"> • Cadmium not detected in fish tissue • Low magnitude of dietary HQs ≥ 1 • Diet assumptions overestimate dietary exposure • Site-specific pathology data did not show abnormalities • Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
White Perch	Chromium	Tissue HQs	<ul style="list-style-type: none"> • WB tissue and dietary HQs < 1 in most scenarios • Tissue and dietary TRVs conservative • Liver HQs superseded by WB tissue HQs • Site-specific pathology data did not show abnormalities • Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
White Perch	Copper	Dietary HQs	<ul style="list-style-type: none"> • Dietary HQs < 1 • Low confidence in the tissue and dietary TRVs • Diet assumptions overestimate dietary exposure • Site-specific pathology data did not show abnormalities • Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Uncertain	Uncertain

Table 11-1a
Weight of Evidence Risk Summary for NBSA BERA

Receptor	Preliminary COC ^a	Primary LOE	Summary of LOE	Risk Conclusions	Risk Driver ^b
White Perch	Mercury	Tissue HQs	<ul style="list-style-type: none"> • WB Tissue LOAEL HQs < 1 • Low magnitude of dietary HQs ≥ 1 • Liver HQs < 1 • Liver HQs superseded by WB tissue HQs • Site-specific pathology data did not show abnormalities • Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
White Perch	Selenium, Silver	Background	<ul style="list-style-type: none"> • Selenium related to background conditions • Tissue and dietary HQs < 1 • Liver HQs ≥ 1 based on unbounded NOAEL • Liver HQs superseded by WB tissue HQs • Site-specific pathology data did not show abnormalities • Long-term fish community monitoring indicate no impairment relative to nearby regional locations 	Unlikely	No
Assessment Endpoint: Survival, growth, and/or reproduction of birds					
Spotted sandpiper	Total Dioxin/Furan/PCB TEQ Bird, Total HMW PAH, Lead, Mercury	Dietary HQs	<ul style="list-style-type: none"> • LOAEL HQs < 1 for most scenarios • Low magnitude of HQs ≥ 1 • Diet and SUF assumptions likely overestimate risk • MDC used as EPC in some cases 	Unlikely	No
Great blue heron	Total Dioxin/Furan/PCB TEQ Bird, Total PCB TEQ Bird, Total DDx (2,4 & 4,4), Mercury, Methylmercury	Dietary HQs	<ul style="list-style-type: none"> • LOAEL HQs < 1 for most scenarios • Low magnitude of HQs ≥ 1 • Diet and SUF assumptions likely overestimate risk • MDC used as EPC in some cases 	Unlikely	No
Double-crested cormorant	Total Dioxin/Furan/PCB TEQ Bird, Total PCB TEQ Bird, Total PCBs	Site- and Species-Specific LOE	<ul style="list-style-type: none"> • Low magnitude of HQs ≥ 1 • NBSA and LPR FFS TRVs are conservative • Site-specific nesting, reproduction, and development study (Parsons 2003) did not show adverse effects 	Unlikely	No
Double-crested cormorant	Total DDx (2,4 & 4,4)	Site-Specific LOE	<ul style="list-style-type: none"> • Low magnitude of dietary and egg HQs ≥ 1 • Site-specific nesting, reproduction, and development study (Parsons 2003) did not show adverse effects 	Unlikely	No
Double-crested cormorant	Mercury, Methylmercury	Dietary HQs	<ul style="list-style-type: none"> • LOAEL HQs < 1 for most scenarios • Low magnitude of dietary and egg HQs ≥ 1 • Site-specific nesting, reproduction, and development study (Parsons 2003) did not show adverse effects 	Unlikely	No

Table 11-1a
Weight of Evidence Risk Summary for NBSA BERA

Receptor	Preliminary COC ^a	Primary LOE	Summary of LOE	Risk Conclusions	Risk Driver ^b
Double-crested cormorant	Copper	Dietary HQs	<ul style="list-style-type: none"> • LOAEL HQs < 1 for most scenarios • Low magnitude of dietary HQs ≥ 1 • Site-specific nesting, reproduction, and development study (Parsons 2003) did not show adverse effects 	Unlikely	No
Lesser scaup	Lead	Dietary HQs	<ul style="list-style-type: none"> • LOAEL HQs < 1 for most scenarios • Low magnitude of HQs ≥ 1 • EPCs for Southwest Zone calculated using clam MDCs 	Unlikely	No
Assessment Endpoint: Survival, growth, and/or reproduction of mammals					
River Otter	2,3,7,8-TCDD, Total Dioxin/Furan TEQ, Dioxin/Furan/PCB TEQ, Total PCBs	Dietary HQs	<ul style="list-style-type: none"> • LOAEL HQs < 1 in most scenarios • Low magnitude of HQs ≥ 1 • SUF assumptions likely overestimate risk 	Unlikely	No
Muskrat	Mercury	Dietary HQs	<ul style="list-style-type: none"> • LOAEL HQs < 1 in most scenarios • Low magnitude of HQs ≥ 1 	Unlikely	No
Harbor Seal	None	Dietary HQs	<ul style="list-style-type: none"> • All HQs < 1 	Not expected	No
Assessment Endpoint: Survival, growth, and/or reproduction of reptiles					
Diamondback terrapin	None	Qualitative Evaluation	<ul style="list-style-type: none"> • Exposures assumed to be similar to or less than for mammals in NBSA • Surface water quality acceptable • Quantitative assessment for some metals in LPRSA (Windward 2019) did not find unacceptable risk 	Not expected	No

Table 11-1a
Weight of Evidence Risk Summary for NBSA BERA

Notes:

^a Preliminary COCs are identified as COPECs/receptors with a LOAEL HQ greater than or equal to 1 for any exposure areas or evaluation scenarios.

^b Risk Drivers identified based on LOAEL HQs ≥ 1 and supporting LOEs.

Acronyms and Abbreviations:

BERA = Baseline Ecological Risk Assessment

COPEC = constituent of potential ecological concern

EPC = exposure point concentration

FFS = focused feasibility study

HQ = hazard quotient

LOAEL = lowest observed adverse effect limit

LOE = line of evidence

LPR = Lower Passaic River

MDC = maximum detected concentration

NOAEL = no observed adverse effect limit

NBSA = Newark Bay Study Area

PCB = polychlorinated biphenyl

SQT = sediment quality triad

SUF = site use factor

TCDD = tetrachlorodibenzo-p-dioxin

TEQ = toxic equivalent

Total DDx = sum of dichlorodiphenyldichloroethylene, dichlorodiphenyldichloroethane, and dichlorodiphenyltrichloroethane

TRV = toxicity reference value

WB = whole-body

-- = not applicable

References:

Iannuzzi, T.J., T.N. Armstrong, E.R. Long, J. Iannuzzi, and D.F. Ludwig. 2008. Sediment quality triad assessment of an industrialized estuary of the northeastern USA. *Environ Monit Assess* 139:257-275.

Bugel S.M., L.A. White, and K.R. Cooper. 2010. Impaired reproductive health of killifish (*Fundulus heteroclitus*) inhabiting Newark Bay, NJ, a chronically contaminated estuary. *Aquatic Toxicol* 96:182-193.

Bugel, S.M., L.A. White, and K.R. Cooper. 2011. Decreased vitellogenin inducibility and 17 β -estradiol levels correlated with reduced egg production in killifish (*Fundulus heteroclitus*) from Newark Bay, New Jersey. *Aquatic Toxicol* 105:1-12.

Bugel, S.M., J.A. Bonventre, L.A. White, R.L. Tanguay, and K.R. Cooper. 2014. Chronic exposure of killifish to a highly polluted environment desensitizes estrogen-responsive reproductive and biomarker genes. *Aquatic Toxicol* 152:222-231.

Parsons, K.C. 2003. Chemical Residues in Cormorants from New York Harbor and Control Location. Contract No. C003858. Submitted to NYSDEC February 28.

Wintermyer, M.L., and K.R. Cooper. 2003. Dioxin/furan and polychlorinated biphenyl concentrations in eastern oyster (*Crassostrea virginica* Gmelin) tissues and the effects on egg fertilization and development. *J Shellfish Res* 22(3):737-746.

Windward Environmental, LLC. 2019. Lower Passaic River Study Area Baseline Ecological Risk Assessment. Final. June.